



Sacch. cerevisiae + *Sacch. cerevisiae ex r.f. bayanus*

ES FLORAL

enartis FERM

YEAST FOR FLORAL WINES

NEW SYNERG-YEASTIC FORMULA

ES FLORAL, yeast selected for its capability to produce intense floral aromas, it is now a blend of strains with enhanced fermentation and sensory characteristics.

SENSORY CHARACTERISTICS

ES FLORAL, a yeast that is distinguished by its capability to produce floral-type aromas, is now a blend of enhanced fermentation and sensory characteristics.

The variety of enzymatic activities present in the different strains make the aroma produced by ES FLORAL more intense and complex. Given a good source of amino acids this blend synthesizes elegant white rose, hawthorn and white fruit notes (peach, apricot, pear etc.).

The palate is pleasantly acidic and fresh.

ES FLORAL can be used for the fermentation of easy-to-drink white, young red and rosé wines.

MICROBIOLOGICAL CHARACTERISTICS

Fermentation temperature	10 - 25°C (50-77°F)
Lag phase	medium
Fermentation speed	moderate
Alcohol tolerance	≤ 15% v/v
Killer factor	neutral
Resistance to free SO ₂	high

ENOLOGICAL CHARACTERISTICS

Nitrogen needs	medium
Oxygen needs	medium-low
Volatile acidity production	low
H ₂ S production	medium
SO ₂ production	low
Glycerol production	medium (6-8 g/L)
Compatibility with malolactic fermentation:	good

APPLICATIONS

Easy-to-drink white, red and rosé wines



Sacch. cerevisiae + *Sacch. cerevisiae ex r.f. bayanus*

ES FLORAL

enartis FERM

MAXIMIZING QUALITY

To enhance fermentation aroma production, it is necessary to ensure adequate yeast nutrition by providing a good source of amino acids and survival factors. Hence, it is advisable to provide **ES FLORAL** with a complex nutrient such as **Nutriferm Arom Plus** at inoculation. **Nutriferm Arom Plus** provides sterols and unsaturated fatty acids useful for maintaining effective cellular metabolism in the presence of alcohol and also amino acids that act as precursors for aromatic compound synthesis.

Using **Enartis Pro Uno** or **Enartis Tan Elegance** during fermentation protects and stabilizes secondary aromas with resulting aromatic longevity in wine.

DOSAGE

20-40 g/hL (1.67 – 3.3 lb/1000 gal)

The highest dosages are recommended in case of rotten grapes, high sugar content and difficult microbiological conditions.

INSTRUCTIONS FOR USE

- Suspend dry yeast in 10 times its weight in clean, warm (35-38°C or 95-100°F) water. Stir gently.
- Let suspension stand for 20 minutes, then stir gently again.
- Add suspension to juice when beginning to fill fermentation tank. The difference in temperature between yeast suspension and juice should not exceed 10°C (18°F).
- Homogenize by pump-over or mixing inoculated juice.

Working to the above-mentioned times and methods ensures maximum activity of re-hydrated yeast.

PACKAGING AND STORAGE

Vacuum packed in 0.5 and 10 kg

Sealed package: store in a cool (preferably 5-15°C or 41-59°F), dry place.

Opened package: carefully reseal and store as indicated above; use quickly.

Product approved for winemaking by the TTB.

Legal Limit: N/A

Product conforms to the *Codex Œnologique International*.

Product approved for winemaking in accordance with
Reg. (EC) N. 606/2009

Contains E 491 Sorbitan monostearate